

AL.2.1992-16

CANADIANA  
DEC 30 1991

**OLDMAN RIVER DAM  
WILDLIFE HABITAT MITIGATION  
- VEGETATION ESTABLISHMENT -**

**Interim Progress Report  
1990-91**

**A. SMRECIU and J. HOBDEN**

Wild Rose Consulting, Inc., Edmonton  
in co-operation with  
Alberta Environmental Centre, Vegreville

Prepared for Alberta Public Works, Supply and Services  
March 1991



Digitized by the Internet Archive  
in 2016

<https://archive.org/details/oldmanriverdamwi00smre>

## TABLE OF CONTENTS

1. INTRODUCTION	1
2. MITIGATION PROJECTS	2
2.1 Auditing Existing Mitigation Projects	2
2.1.1 Shallow Water	2
2.1.2 Deep Water	2
3. OLDMAN RIVER DAM	3
4. WILDLIFE HABITAT MITIGATION	4
5. - VEGETATION ESTABLISHMENT -	5
6. SPECIES DOCUMENTATION	6
6.1 Species Documentation	6
6.2 Seed Bank	6
6.2.1 Seed Production	6
6.2.2 Seed Collection	6
7. REFERENCES CITED	7

**A. SMRECIU and J. HOBDEN**

Wild Rose Consulting, Inc., Edmonton  
in co-operation with  
Alberta Environmental Centre, Vegreville

Prepared for Alberta Public Works, Supply and Services  
March 1991

OLDMAN RIVER DAM  
WILDLIFE HABITAT MITIGATION  
VEGETATION ESTABLISHMENT

Initial Progress Report  
1990-91

A. GEMMILL and J. HODGE

Wild Area Consulting, Inc., Edmonton  
in co-operation with  
Alberta Environmental Centre, Vegreville

Prepared for Alberta Public Works, Supply and Services  
March 1991



## TABLE OF CONTENTS

1 INTRODUCTION .....	1
2 MITIGATION PROJECTS .....	5
2.1 Auditing Existing Mitigation Projects .....	5
2.1.1 Shelterbelt and nursery plantings .....	5
2.1.2 Snow fence effects on tree and shrub encroachment .....	5
2.1.3 Soil moisture behind snow fences .....	8
2.1.4 Grassland recovery .....	13
2.1.5 Grassland restoration .....	17
3 SPECIES DOCUMENTATION AND SEED BANK .....	28
3.1 Species Documentation .....	28
3.2 Seed Bank .....	29
3.2.1 Seed Production .....	29
3.2.2 Seed Collection .....	30
4 REFERENCES CITED .....	31

# TABLE OF CONTENTS

1	INTRODUCTION	1
2	MITIGATION PROJECTS	2
2.1	Auditing Existing Mitigation Projects	2
2.1.1	Shallow and water planting	2
2.1.2	Shore bank erosion on low and high water	2
2.1.3	Soil erosion control and riprap	2
2.1.4	Grassland recovery	2
2.1.5	Grassland restoration	2
3	RESEARCH DOCUMENTATION AND DATA BASE	3
3.1	Species Documentation	3
3.2	Soil Bank	3
3.2.1	Soil Production	3
3.2.2	Soil Collection	3
4	REFERENCES CITED	4

## LIST OF TABLES

<b>Table 1.</b> Vegetation cover in plots where snow fence was placed to encourage invasion of shrubs from coulee slopes into surrounding grassland. ....	7
<b>Table 2.</b> Species abundance in grassland recovery plots. ....	14
<b>Table 3.</b> Average total productivity in 1989 and 1990. ....	23
<b>Table 4.</b> Effect of direction from fence line on productivity. ....	23
<b>Table 5.</b> Effect of direction from the fence line on grass productivity. ....	23
<b>Table 6.</b> Effect of snow fence on grass productivity windward of the fence line. ....	23
<b>Table 7.</b> Mean squares (MS) of each of the primary effects for soil moisture analysis at each of four depths. ....	25

# LIST OF TABLES

Table 1. Frequency of occurrence of plants whose names were placed in botanical literature of America from 1800 to 1850 (including present)	1
Table 2. Species abundance in botanical literature of America	14
Table 3. Species abundance in botanical literature of America in 1800 and 1850	23
Table 4. Effect of climate on species abundance in botanical literature of America	23
Table 5. Effect of climate on species abundance in botanical literature of America in 1800 and 1850	23
Table 6. Effect of climate on species abundance in botanical literature of America in 1800 and 1850	23
Table 7. Effect of climate on species abundance in botanical literature of America in 1800 and 1850	23



## LIST OF FIGURES

<b>Figure 1.</b> Location of study plots at the Oldman River Dam Site. ....	4
<b>Figure 2.</b> Effects of snow fences on soil moistures (at 0-15 cm) in mid-summer and fall at seven distances from the fence line. ....	11
<b>Figure 3.</b> Effects of snow fences on soil moistures (at 30-45 cm) in mid-summer and fall at seven distances from the fence line. ....	11
<b>Figure 4.</b> Effects of snow fences on soil moistures (at 60-75 cm) in mid-summer and fall at seven distances from the fence line. ....	12
<b>Figure 5.</b> Effects of snow fences on soil moistures (at 90-105 cm) in mid-summer and fall at seven distances from the fence line. ....	12
<b>Figure 6.</b> Field design of experiment to determine effects of snow fence on soil moisture and above-ground biomass. ....	22
<b>Figure 7.</b> Effects of snow fence on total above-ground biomass at five distances in each direction (windward and leeward) from the fence line. ....	24
<b>Figure 8.</b> Effects of snow fence on above-ground biomass of grasses at five distances in each direction (windward and leeward) from the fence line. ....	24
<b>Figure 9.</b> Effects of snow fences on fall soil moisture at 0-15 cm, at seven distances, leeward and windward of the fence line. ....	26
<b>Figure 10.</b> Effects of snow fences on fall soil moisture at 30-45 cm, at seven distances, leeward and windward of the fence line. ....	26
<b>Figure 11.</b> Effects of snow fences on fall soil moisture at 60-75 cm, at seven distances, leeward and windward of the fence line. ....	27
<b>Figure 12.</b> Effects of snow fences on fall soil moisture at 90-105 cm, at seven distances, leeward and windward of the fence line. ....	27

# LIST OF FIGURES

Figure 1. Location of study plots at the Oltman River near the	1
Figure 2. Effects of snow fences on soil moisture at 0-15 cm in mid-winter and fall	42
Figure 3. Effects of snow fences on soil moisture at 30-45 cm in mid-winter and fall	44
Figure 4. Effects of snow fences on soil moisture at 60-75 cm in mid-winter and fall	46
Figure 5. Effects of snow fences on soil moisture at 90-105 cm in mid-winter and fall	48
Figure 6. Field design of experiment to determine effects of snow fences on soil moisture	50
Figure 7. Effects of snow fences on total above-ground biomass at five distances in each	52
Figure 8. Effects of snow fences on above-ground biomass of grasses at five distances in	54
Figure 9. Effects of snow fences on fall soil moisture at 0-15 cm at seven distances	56
Figure 10. Effects of snow fences on fall soil moisture at 30-45 cm at seven distances	58
Figure 11. Effects of snow fences on fall soil moisture at 60-75 cm at seven distances	60
Figure 12. Effects of snow fences on fall soil moisture at 90-105 cm at seven distances	62

## LIST OF APPENDICES

<b>Appendix I.</b> Climate information for Pincher Creek (1989-1990) .....	33
<b>Appendix II.</b> Vascular Plants of the Oldman River Dam Site. ....	34
<b>Appendix III.</b> Voucher specimens collected .....	41
<b>Appendix IV.</b> Native seed collection .....	49





## **ACKNOWLEDGEMENTS**

The authors are grateful to Al Nilson of Alberta Public Works, Supply and Services for his guidance and enthusiasm throughout the year, and to June Carrington and Craig Osterloh for keeping us up to date on progress at various mitigation projects.

We would like to acknowledge the logistical support of the Alberta Environmental Centre (Vegreville). A special thanks to Reinhard Hermesh and Barbara Darroch for their valuable input, ongoing support, and continued patience. We are grateful to Zack Florence for his expert advice concerning experimental designs and statistical analyses. Thanks to Denise Burton and Leslee Porayko for their technical assistance.

Thanks to Alberta Environment (especially Mr. Dale Hay) for installation of the moisture tubes.

We are extremely grateful to Mr. Dave Hay for preparing and assisting in field arrangements.



# 1 INTRODUCTION

The Oldman River Dam, first announced in 1980, is being built in southwestern Alberta 10 km northeast of Pincher Creek, just east of the confluence of the Oldman, Castle, and Crowsnest Rivers (Figure 1), a site confirmed in 1984. The primary purpose of the dam and the resulting reservoir will be flow regulation and replenishing storage during maximum flow (mid-May to mid-July) to provide a dependable water supply throughout the year for domestic, municipal, industrial, agricultural, and recreational uses. Construction of the project began in 1986 and is expected to be complete in late 1991. Water was first stored in the spring of 1991. At full supply level, the reservoir will cover an estimated area of 2400 ha, and will be approximately 24 km long and up to 3 km wide. Water depths will exceed 60 m in some parts.

To cope with the environmental disturbances that would inevitably arise from the construction and eventual flooding, the Planning Division of Alberta Environment initiated studies which lead to Environmental Mitigation/Opportunities Plans for various aspects of concern including environment (wildlife and fisheries), historical resources, and recreation (Alberta Environment 1988a, 1988b).

Work began on the wildlife portion of the Environmental Mitigation/Opportunities Plan in 1985 and 1986 with a review of pertinent information, followed by field surveys of wildlife. A vegetation survey, conducted in 1984 and 1985, described major plant communities and discussed dominant factors controlling vegetation in the areas surrounding the future reservoir site (Hardy Associates (1978) Ltd. 1986). The report also included a brief reclamation plan and recommendations concerning procedures and methods for preserving and restoring native vegetation.

The Delta Environmental Management Group Ltd. (the Delta Group) was retained by Alberta Environment in 1987 to prepare a strategy for wildlife habitat mitigation at the Oldman River Dam Site. Their objectives were to:

'...identify opportunities for wildlife habitat mitigation in the vicinity of the reservoir and, on the basis of this inventory, develop practical and effective projects to protect, enhance, or create wildlife habitat during 1988 to 1993.' (Green and Eccles 1989).

## Oldman River Dam Wildlife Habitat Mitigation - Vegetation Establishment

An action plan was formulated by the Delta Group in cooperation with Alberta Public Works, Supply and Services (APWSS) (Nilson and Green 1989). Projects were presented along with an implementation plan for wildlife habitat mitigation projects.

Three major mitigation techniques were recommended and are now being implemented. The first is **protection** of important wildlife habitat which already exists in the vicinity of the reservoir. This is being accomplished by acquiring both crown and private lands. The land base has been fenced to exclude grazing and other activities that would be detrimental to the native habitat. The second major method is habitat **enhancement**. This includes techniques undertaken to modify or improve sites to increase their use by wildlife. Enhancement included placement of snow fences to improve soil moisture and planting of trees and shrubs around existing wetlands and in other suitable areas. New wildlife habitats are also being **created** by construction of riparian dams and check dams and planting appropriate plant species (i.e. those which will provide erosion control and provide food and shelter for wildlife). Other projects involve creating or enhancing nesting sites for raptors, providing suitable structures for duck nesting sites and providing structures for other animals. These however, do not involve modifying vegetation and therefore will not be considered further in this report.

Since the objective of wildlife habitat mitigation is to provide wildlife with sufficient habitat to replace that being lost to flooding after 1991, the success of the program will be measured by the eventual use by wildlife and survival of wildlife after flooding. Success of many individual projects however, depends on establishment of specific (mostly native) plants and enhancement of specific vegetation. Success also depends on the potential for natural regeneration of improved vegetation. Only the vegetation aspects of specific projects are considered in results reported here.


Beginning in June of 1989, various mitigation projects at the Oldman River Dam site were surveyed and audited. Experimental programs were set up, in co-operation with Plant Sciences Division (Vegetation Branch) of Alberta Environmental Centre (AEC) in Vegreville, to evaluate and where necessary improve the effectiveness of existing methods of protecting and enhancing plant communities, to evaluate methods and species selection in the creation of new wildlife habitat, and to provide input into planning for supplemental and replacement plantings.



An Interim Progress Report was submitted to APWSS in March 1990 (Smreciu 1990). That report described the specific background, plans and experimental designs for projects which were started prior to 1989, and outlined those to be started in 1990.

This interim report consists of results from surveys of projects which were started prior to 1989, and the design and preliminary results of experiments started in 1990. Also included is a preliminary list of vascular plant species native to the area of the dam and reservoir and lists of voucher specimens and seed collected in 1989 and 1990. Scientific names used in this report follow Moss (1983), Smreciu and Currah (1989), and Aiken and Darbyshire (1990).

Figure 1. Location of study plots at the Oldman River Dam Site.



## 2 MITIGATION PROJECTS

The following is a brief description of the monitoring and experimental projects which have been set up in the land base around the Oldman River Dam site and presents the interim results from the monitoring of each in 1990. Figure 1 illustrates the location of each project on-site.

### 2.1 Auditing Existing Mitigation Projects

#### 2.1.1 Shelterbelt and nursery plantings

**Background:** In 1987 and 1988 a major tree planting scheme was undertaken on a site east of the dam (PROJECTS OS26 (Sec 21, Twp 7, Rg 29, W4M) and OS17 (Sec 16, Twp 7, Rg 29, W4M)). Numerous snow fences were erected and trees were planted on the leeward sides and supplied with moisture through a drip irrigation system. Most of the site had been previously cultivated and has been seeded to native and introduced forage grasses. Small areas closer to the river valley had not been cultivated and trees planted in this area were planted in narrow trenches cut into the native grassland. Shelterbelt plantings were placed to reduce soil erosion and provide shelter for wildlife. Survival of trees was not recorded in 1987 and 1988. In 1989, 10 plots were designated and the number of live and dead trees were counted in each plot. Results in 1989 indicated that most of the trees were still alive but that the Colorado blue spruce (*Picea pungens* 'glauca') and the Douglas fir (*Pseudotsuga menziesii*) were in poor shape and continued survival was doubtful. The demise of these trees was probably due to winter desiccation and the alkalinity of the soil. The caragana (*C. arborescens*) had survived from 1988 when many had been replanted, however severe insect damage had occurred especially on those planted into the native range or in areas where the introduced forage grasses had become established. Northwest poplar had survived but some damage had been caused by deer browsing.

Monitoring survival and growth of these trees is being done every two years therefore no monitoring was undertaken at this site in 1990. Monitoring will continue in 1991.

#### 2.1.2 Snow fence effects on tree and shrub encroachment

**Background:** Coulees, on the north side of the future reservoir and oriented in a north-south direction, have limited numbers of trees and shrubs, most of which grow on the east face where they are protected from wind, and in the bottom of the coulees where moisture collects. At the top edge of the coulees shrub communities are sharply differentiated from the surrounding grassland

community because there is no longer protection from the drying winds and soil moisture is reduced. Since shrubs and trees provide better cover and browse for wildlife than grasslands, snow fences were erected in the fall of 1988 (by APWSS) at a site to the north and east of the dam, within the protected land base (PROJECT OS17 (Sec 16, Twp 7, Rg 29, W4M)). Snow fences were placed along the ridges between the coulees to encourage the encroachment of trees and shrubs from the coulee up into the grassland. In 1989 five transects were set up from a NW-SE fence line to the edge of the shrub communities. Two were placed where there was snow fence, two were in an area with no snow fence and one was placed in a transition area along each transect. A permanent plot was placed at the edge of the shrub community and three more plots were set up at 2 m intervals from the first. The objective of this audit was to monitor shrub encroachment from the coulee slopes to the adjacent grassland when snow fences are placed along the top of the coulee slope. The relative abundance of vascular plant species will be determined each year and compared to determine if the area covered by shrubs is increasing. Records were taken in 1989 as a baseline.

**Results (1990):** Data collected in 1990 are presented in Table 1. Observations taken in 1990 indicate a marked increase in the relative abundance of both grasses and forbs in most of the plots (with the exception of the forbs in plots of Transect E). Much of this increase is likely due to the differences in precipitation (in amounts and the timing of the rainfall) between 1989 and 1990 (Appendix 1). Some of the increase is due to the protection offered by the snow fences. The decrease in forbs in Transect E plots could be a result of the increase in grass cover.

After only one year there is little indication of increasing shrub cover. Shrub encroachment is a long term effect and little change is expected over a period of one year.



**Table 1.** Vegetation cover in plots set up to determine if snow fence placement increases invasion of shrubs from coulee slopes into surrounding grassland.

Plot Number*	Shrubs (%)		Graminoids (%)		Forbs (%)	
	1989	1990	1989	1990	1989	1990
A-15m	0.0	0.0	26.0	45.0	5.0	18.0
A-17m	0.0	0.0	26.0	31.0	2.0	10.0
A-19m	3.0	10.0	19.0	40.5	3.0	6.0
A-21m	41.0	36.0	2.0	28.0	57.0	52.0
B-20m	0.0	0.0	9.0	20.0	8.0	15.0
B-22m	0.0	0.0	37.0	31.0	3.0	16.0
B-24m	18.0	40.0	21.0	45.0	6.0	18.5
B-26m	35.0	43.0	20.0	25.0	8.0	15.0
C-21m	0.0	0.0	11.0	33.0	16.0	18.0
C-23m	0.0	0.0	11.0	46.0	9.0	21.0
C-25m	0.0	0.0	6.0	15.0	22.0	37.5
C-27m	55.0	67.5	30.0	36.0	7.0	12.5
D-22m	0.0	0.0	35.0	36.0	17.0	19.5
D-24m	0.0	0.0	29.0	35.0	16.0	20.5
D-26m	0.0	0.0	45.0	31.5	15.0	11.5
D-28m	46.0	40.0	31.0	41.5	16.0	24.5
E-22m	0.0	0.0	12.0	35.0	25.0	11.0
E-24m	0.0	0.0	4.0	30.0	35.0	9.5
E-26m	0.0	0.0	9.0	53.0	36.0	9.0
E-28m	28.0	18.0	27.0	41.5	41.0	10.5

\* A and B are snow fenced, C is in the transition zone and D and E are from areas with no snow fence. Distances indicated in the plot numbers are from the fence line.

### 2.1.3 Soil moisture behind snow fences

**Background:** Snow fencing was erected on a section of perimeter fence at a site west of the Castle River and south of Highway 3 (PROJECT C37 (Sec 11, Twp 7, Rg 1, W5M)) in the fall of 1988. The primary purpose was to increase soil moisture levels such that trees and shrubs planted as part of the wildlife habitat mitigation program would have an increased chance of survival. Snow fences were placed at intervals, and gaps were left to allow access to wildlife. The existing fence provided an opportunity to study the effects of snow fences on soil moisture and provide site specific information for planning tree and shrub placement in the future. An experiment was set up to determine if significant differences in soil moisture could be detected among areas behind the snow fences and the controls (areas with no snow fences). Comparisons were made for various distances from the fence (1 m, 2 m, 4 m, 8 m, 16 m, and 32 m), at various depths (15 cm, 30 cm, 45 cm, 60 cm, and 75 cm) and at two different times in the season (summer and fall). A hand auger was used to obtain soil samples at the various depths. The percentage soil moisture (by weight) was calculated.

First year results indicate that there was a significant difference in soil moisture between fenced and unfenced areas and that depth, distance, and season were also significant factors. Generally soils were drier at greater depths, although surface samples were drier than those taken just below surface. Soils farther from the fence line were moister due to water run-off down slope, away from the fenceline. The greatest increases in soil moisture due to snow fences were at 2 m, 4 m, and 8 m from the fence (Smreciu 1990).

Moisture tubes were inserted in the ground in early summer of 1990 and soil moisture measurements were made with a Campbell Pacific (503) Moisture Gauge twice in the season (summer and fall) at depths of 0-15 cm, 30-45 cm, 60-75 cm, and 90-105 cm. Soil moisture results in 1990 were percentages by volume.

**Results (1990):** The surface soil moisture (depth 15 cm) was highest in the summer, behind the snow fences (Figure 2). At this depth the summer controls were higher than the fall snow fenced areas, which in turn were higher than the fall controls. Season had a greater influence on the soil moisture at 0-15 cm than did the effect of the snow fences. The effect of season is a reflection of the higher precipitation in July and August (summer) than in September (fall) (76.6 mm, 33.6

mm, and 10.2 mm respectively) (Appendix 1). Little change in soil moisture was recorded among distances at this depth.

At a depth of 30-45 cm results were similar, although overall moisture was somewhat higher (Figure 3). Down slope (at 24-32 m) in both summer and fall, the soil moisture of the controls slightly surpassed the snow fenced plots. This is probably an indication that the treatments were not independent (i.e. the effects of the snow fences placed on either side of the controls were influencing the controls) and soil moisture increased due to both slope and to snow catchment.

At 60-75 cm the snow fenced results for the fall surpassed those of the summer controls up to approximately 14 m from the fence line (Figure 4). Snow fences improved moisture to a great extent at this depth up to 14 m and the moisture was retained to some extent. At 90-105 cm, again soil moisture measured behind the fences surpassed those recorded in the controls in both summer and fall (Figure 5).

Although in 1989 soils were drier at greater depths (to 75 cm), in 1990 the trend reversed; the recorded soil moisture increased at greater depths to 105 cm especially those behind the snow fences, indicating a retention of soil moisture. A smaller increase was observed at greater depths in the controls.


Generally there were no differences in soil moisture among sampling distances. This is largely due to the slope on which these plots are located. There were greater differences between control and snow fenced sample sites closer to the fences. Moisture at 16 m and 32 m are not as affected by snow fences to the same degree as those at 1 m, 2 m, 4 m, and 8 m. This was largely because the gaps in the snow fences were small (approximately 13 m) and the effects of the snow fences fan out and overlap.

Measurements of soil moisture taken in established shrub communities at seven sites around the Oldman River Dam indicate that the moisture requirement for shrub communities is variable. Fall moisture readings in the shrub communities averaged 16% at 0-15 cm, 22% at 30-45 cm, 27% at 60-75 cm, and 30% at 90-105 cm. These moisture readings were not significantly different from those taken at seven grassland locations around the reservoir. The moisture measured in soils at Project C37 were comparable to those taken in native shrub populations such as saskatoon/choke


cherry, and Douglas fir around the reservoir in the summer and fall indicating there was sufficient moisture during these times to sustain native shrub communities. Spring soil moisture measurements and comparisons would be useful in determining if this area could sustain a shrub community. There are also other factors such as protection from wind, and competition with other plants which would have an effect on shrub establishment in this (and other) areas.




**Figure 2.** Effects of snow fences on soil moistures (at 0-15 cm) in mid-summer and fall at seven distances from the fence line.




**Figure 3.** Effects of snow fences on soil moistures (at 30-45 cm) in mid-summer and fall at seven distances from the fence line.



**Figure 4.** Effects of snow fences on soil moistures (at 60-75 cm) in mid-summer and fall at seven distances from the fence line.



**Figure 5.** Effects of snow fences on soil moistures (at 90-105 cm) in mid-summer and fall at seven distances from the fence line.



#### 2.1.4 Grassland recovery

**Background:** To improve the range for wildlife an area of previously grazed native rangeland was protected from grazing beginning in 1989 (PROJECT CR17 (Sec 32, Twp 7, Rg 1, W5M). In spring of 1989 six plots (each 2 m x 2 m) were placed in the protected area. These will be monitored over a period of at least three years for changes in species composition and relative abundance of individual species. Preliminary data were taken in 1989.

**Results (1990):** Relative abundance was recorded for each vascular plant species in each of the six plots. Results are presented in Table 2. One of the differences from 1989 to 1990 was an increase in the graminoid cover and an increase in the number of grasses observed in many of the plots. There was little change in the shrub cover. The elimination of grazing gave the grasses a short-term, competitive advantage. Due to the growth habit of shrubs, changes in shrub cover will only be observed after several years. There were some differences in the forbs which were observed in 1990. In plots 2, 3, and 6 there was an increase in the forb cover, and overall there were several species observed in 1990 which were not in 1989. Some differences observed between 1988 and 1990 were likely due to changes in precipitation (Appendix 1). Continued monitoring will be required prior to analysing data for compositional and abundance differences.

Table 2. Species cover in grassland recovery plots.

Taxa	Cover					
	Plot 1 (%)	Plot 2 (%)	Plot 3 (%)	Plot 4 (%)	Plot 5 (%)	Plot 6 (%)
<b>Shrubs</b>						
<i>Rosa arkansana</i>	1	t*	5	2	1	0
<i>Symphoricarpos occidentalis</i>	0	0	17	0	0	0
<b>Graminoids</b>						
<i>Agropyron smithii</i>	1	10	1	5	25	25
<i>Agropyron spicatum</i>	5	0	0	0	1	0
<i>Agropyron subsecundum</i>	0	0	1	0	0	0
<i>Carex</i> sp.	0	2	0	5	3	1
<i>Danthonia parryi</i>	15	5	20	t	t	t
<i>Festuca campestris</i>	10	t	15	1	0	1
<i>Koeleria macrantha</i>	15	20	3	30	10	15
<i>Stipa curtisetia</i>	2	1	t	15	2	1
<b>Forbs</b>						
<i>Achillea millefolium</i>	t	0	2	0	0	5
<i>Agoseris glauca</i>	t	0	0	0	0	0
<i>Allium cernuum</i>	0	t	1	0	0	0
<i>Allium textile</i>	0	0	t	0	0	0
<i>Anemone multifida</i>	0	0	t	0	0	0
<i>Antennaria parvifolia</i>	20	5	0	8	5	5
<i>Artemisia campestris</i>	t	0	t	0	t	0
<i>Artemisia frigida</i>	10	10	4	5	10	8
<i>Aster ericoides</i>	1	5	8	5	4	3
<i>Aster laevis</i>	1	0	0	0	0	0
<i>Astragalus flexuosus</i>	0	0	t	t	0	5

\* t = &lt; 1%.



Table 2. Species cover in grassland recovery plots.

Taxa	Cover					
	Plot 1 (%)	Plot 2 (%)	Plot 3 (%)	Plot 4 (%)	Plot 5 (%)	Plot 6 (%)
<i>Astragalus dasyglottis</i>	t*	0	0	0	0	0
<i>Cerastium arvense</i>	2	0	1	0	0	0
<i>Cirsium undulatum</i>	0	t	0	0	0	0
<i>Comandra umbellata</i>	0	t	0	0	t	t
<i>Cryptantha nubigena</i>	0	1	0	0	0	0
<i>Erigeron caespitosus</i>	1	t	t	0	0	0
<i>Gaillardia aristata</i>	t	t	1	t	1	t
<i>Galium boreale</i>	2	0	2	0	3	t
<i>Geum triflorum</i>	0	3	5	0	0	0
<i>Heterotheca villosa</i>	t	2	0	0	0	0
<i>Heuchera parvifolia</i>	0	0	0	t	0	0
<i>Liatris punctata</i>	2	3	0	t	3	0
<i>Lithospermum ruderales</i>	5	0	2	t	10	5
<i>Orthocarpus luteus</i>	t	0	5	0	t	1
<i>Oxytropis monticola</i>	1	0	0	0	0	0
<i>Oxytropis sericea</i>	1	t	t	t	0	0
<i>Penstemon procerus</i>	0	0	t	0	0	0
<i>Petalostemon purpureum</i>	1	1	0	0	0	0
<i>Phlox hoodii</i>	1	3	5	8	2	0
<i>Potentilla concinna</i>	0	t	0	0	0	0
<i>Potentilla hippiana</i>	t	t	t	0	0	0
<i>Potentilla pensylvanica</i>	t	0	t	t	0	t
<i>Senecio canus</i>	0	0	5	0	5	0
<i>Solidago missouriensis</i>	0	0	5	0	5	0

\* t = < 1%

**Table 2.** Species cover in grassland recovery plots.

	Cover					
	Plot 1 (%)	Plot 2 (%)	Plot 3 (%)	Plot 4 (%)	Plot 5 (%)	Plot 6 (%)
<i>Sphaeralcea coccinea</i>	0	0	0	0	0	1
<i>Vicia americana</i>	t*	0	0	t	0	0
<b>Ruderals</b>						
<i>Erysimum inconspicuum</i>	0	0	0	t	1	t
<i>Taraxacum officinale</i>	t	0	0	0	0	0
<i>Tragapogon dubius</i>	0	t	0	0	0	0

\* t = < 1%

### 2.1.5 Grassland restoration

**Background:** An experiment was designed and set up in 1989 to study the effects of snow fences on the recovery of heavily grazed grassland. Although snow fencing was the primary factor being studied the experiment was designed to look at several other factors including direction from fences (windward/leeward), distance from the fences (2 m, 4 m, 8 m, 16 m, and 32 m) and their effects on the above-ground biomass of vascular plant species over six years. The field design of the experiment was presented in Smreciu (1990) and a diagram is included here for reference (Figure 6).

In 1989, prior to snow fence placement, above-ground biomass data were gathered and analysed. No statistical differences in total above-ground biomass were apparent among the plots. Snow fences were placed in October, 1989.

In 1990 biomass was cut in late August and early September and dried until mid-September when dry weights were recorded. Results were analysed and compared to those collected in 1989. Relative abundance and species composition were recorded for each vascular plant species in one sub-plot of each of the control and snow fenced plots in both 1989 and 1990.

In 1990 a second experiment was added onto the original design. This second experiment was set up to study the effects of snow fences on soil moisture at the Oldman River Dam Site. In early summer, soil moisture tubes were placed to one side or the other of the strips of plots and also at distances of 12 m and 24 m. Soil moisture was measured (using a Campbell Pacific (503) Moisture Gauge) in mid-summer and in the fall. Data were analysed using a GLM and an LSD test (SAS Institute 1988).

**Results (1990):** As expected there were significant differences between above-ground biomass data collected in 1989 and those recorded in 1990 (Table 3). There was a significant increase in total biomass throughout the entire experiment due in part to the differences in precipitation recorded in the region between 1989 and 1990 (i.e. there was greater rainfall during May, a time of active growth for many plants, in 1990 then in 1989) (Appendix 1). The remaining differences were attributed to the elimination of grazing from this area in 1990 and the placement of snow fences in late 1989 after baseline data had been collected.

Since there were highly significant differences in biomass from 1989 and 1990 these were analysed separately. As reported last year there were no differences among the different treatments in 1989 (Smreciu 1990). This was to be expected since no treatments had been started when the initial biomass was clipped.

Biomass data collected in 1990 were analysed and a highly significant difference was observed between data collected from the windward side of the fence line and those collected from the leeward side of the fence line (Table 4). Further analysis of data from the windward plots indicated that differences between snow fenced and control plots were not significant, and although differences between the biomass at different distances varied, no significance was found among that data either.

Analysis of above-ground biomass collected from leeward plots indicated that neither distance nor snow fences were significant factors, but there was a significant interaction between the two factors (i.e. at some distances the snow fences had an influence on the amount of biomass). When the means for the snow fenced plots and the control plots were plotted against each other at each of the distances (Figure 7) we see that the reason for the interaction is that the treatments were not independent. The effect of snow fence seems to be wider as distances from the fence line increased and there is an area of overlap beyond 16 m.

Where snow fences were placed there was a peak biomass at approximately 4 m from the fence line and this dropped off to 32 m. Although differences in biomass among the various distances were not significant, when monitored for several years these differences might become more pronounced. In the control plots (those where no snow fences were placed) the biomass is low at the 2-4 m distances but increases toward the 16 m distance and then levels out somewhat toward 32 m from the fences. The increase in the biomass at 16 m indicates the overlap of effects from the fenced areas as mentioned above.

An interesting drop in biomass is apparent at 2 m on the windward side of the fence line. This reduction in biomass is probably due to damage caused by the construction of the fence (i.e. vehicle movement along the fence line). It is interesting to note that the effect is much reduced where snow fences are in place.



Biomass data for grasses only were similar to those for the total biomass (Figure 8). Windward and leeward results for grasses were significantly different from each other (Table 5).

Whereas on the windward side of the snow fence there was no significant difference in **total biomass** between fenced and control plots there was a significant difference in **grass biomass** between fenced and control plots on the windward side (Table 6). Differences among distances (windward, fenced) were not significant.

Leeward of the fence line there was greater grass biomass than on the windward side however, no significant differences were found between fenced and control plots. As with the windward results there was some variation in the amount of herbage recorded at different distances but these had no statistical significance (Figure 8).

A drop in grass biomass, similar to that observed in the total biomass results, was recorded at 2 m from the fence on the windward side but **only** in the controls. This suggests that the grasses were able to recover more quickly than the forbs and shrubs but only when sufficient moisture was available (see discussion below regarding soil moisture responses to the snow fences).

No significant differences were observed for the biomass data for forbs or shrubs. These plants comprised a smaller portion of the vegetation and therefore relatively large changes due to snow fence placement would have to be recorded before results would be significant. These plants also take longer than grasses to respond to changes in soil moisture because of their growth habits. Forbs have long root systems which absorb moisture from greater depths than grasses and are somewhat more determinant in the number of shoots which can be produced as a response to moisture in a single year. Shrubs, because of their woody habit require a longer period to increase their above-ground component. The biomass of each of these groups is not tied to annual moisture cycles to the same extent as is the grass biomass. Perhaps over time differences among these groups will be observed due to the effects of the snow fences.

Soil moisture measurements taken at this site were analysed at each of four depths (0-15 cm, 30-45 cm, 60-75 cm, and 90-105 cm). The primary effect on soil moisture at 0-15 cm was season, and the effect of season diminished at greater depths; at 90-105 cm seasonal effects were not

significant. Season had a significant effect on soil moisture at 30-45 cm and 60-75 cm although the effect of snow fence was greater. As depth increased the effect of season diminished and the effect of the snow fences increased (Table 7). The effects of direction and distance also increased at greater depths. The great changes in soil moisture observed between summer and fall at 0-15 cm (Figure 9) were due to the effect of local precipitation, temperature, and drying winds. Soil moisture recorded at 90-105 cm did not change from summer to fall because there was no influence at this depth from the local daily and monthly climate changes.

Since our primary interest lies in the change in soil moisture due to snow fences (not seasonal changes) the results for the 0-15 cm depth are disregarded in these discussions. The summer moisture was greater than the fall readings at depths of 30-45 cm and 60-75 cm and remained constant at the 90-105 cm depth. We are interested especially in how much moisture is retained in the soil and therefore the fall data were analyzed in greater detail.

When only the fall data were analysed, both snow fence and direction were found to be significant factors at all three depths. The snow fence and direction effects increased with increased depth and at 90-105 cm distance and interactions of distance x fence and distance x direction were significant. Since direction was highly significant in the analysis of the fall data (at each of the three depths), data for each direction (windward and leeward) were analysed separately. Figures 10-12 are graphs comparing fall soil moisture data from the controls with the snow fenced data at each depth (30-45 cm, 60-75 cm, and 90-105 cm).

Analysis of the leeward data at 30-45 cm showed that the snow fence had a significant effect on soil moisture and at 60-75 cm and 90-105 cm the effect was highly significant. Analysis of leeward controls (areas with no snow fences) indicated there were no differences in soil moisture among various distances from the fence line at any of the three depths. For the areas where fences were placed, soil moisture did not vary significantly among distances at 30-45 cm or 60-75 cm leeward of the fence line, but it did vary significantly among distances at 90-105 cm.

Windward of the fence line, snow fences increased soil moisture significantly only at 60-75 cm and 90-105 cm. At 90-105 cm there was also a significant interaction between fence and distance (the snow fences had an effect on soil moisture at some distances).

There was no significance among distances for the controls (windward) at any of the three depths. There were however, significant differences among distances for all three depths, for the snow fenced results. At 30-45 cm the highest moisture were measured at 2 m and 4 m. This was also the case for depths of 60-75 cm and 90-105 cm (Figures 10-12).


In summary, the soil moisture (1990) was greater on the leeward side than on the windward side of the fence line. On the windward side soil moisture was significantly greater in the snow fenced areas than in the controls at depths of 60-75 cm and 90-105 cm. In the control plots, no significant differences were observed among various distances from the fence line, but in the snow fenced plots there were significant differences among distances from the fence line at all three depths. The greatest moisture were observed at 2-4 m from the fence line. Leeward of the fence line, significant differences were observed between control and snow fenced plots. There were no significant differences in soil moisture among distances for the controls and among snow fenced areas, distance was only significant at the 90-105 cm depth.

Greatest soil moistures were recorded at 90-105 cm at distances of 2-8 m behind the snow fences. A similar trend was apparent for the 60-75 cm and 30-45 cm depths but the actual moisture percentages were lower in each case.

In both soil moisture experiments we found that the moisture which is accumulating behind the snow fences is being retained at the 90-105 cm depth. At both sites we found that the accumulated soil moisture behind the snow fences compares favourably with soil moisture measured in shrub communities around the Oldman River Dam site (see page 9).



**Figure 6.** Field design of experiment to determine effects of snowfence on soil moisture and above-ground biomass.





**Table 3.** Average total productivity in 1989 and 1990.

<b>Year</b>	<b>Above-ground biomass<sup>*</sup> (g)</b>
1989	70 a
1990	146 b

<sup>\*</sup> Means followed by the same letter are not significantly different (LSD).

**Table 4.** Effect of direction from fence line on productivity (1990).

<b>Direction</b>	<b>Above-ground biomass<sup>*</sup> (g)</b>
Leeward	158 a
Windward	135 b

<sup>\*</sup> Means followed by the same letter are not significantly different (LSD).

**Table 5.** Effect of direction from the fence line on grass productivity (1990).

<b>Direction</b>	<b>Above-ground biomass<sup>*</sup> (g)</b>
Leeward	117 a
Windward	103 b


<sup>\*</sup> Means followed by the same letter are not significantly different (LSD).

**Table 6.** Effect of snow fence on grass productivity windward of the fence line (1990).


<b>Treatment</b>	<b>Above-ground biomass<sup>*</sup> (g)</b>
Fenced	110 a
Control	95 b

<sup>\*</sup> Means followed by the same letter are not significantly different (LSD).

**Figure 7.** Effects of snow fence on total above-ground biomass at five distances in each direction (windward and leeward) from the fence line.



**Figure 8.** Effects of snow fence on above-ground biomass of grasses at five distances in each direction (windward and leeward) from the fence line.




**Table 7.** Mean squares (MS) of each of the primary effects for soil moisture analysis at each of four depths.


Source	DF	MS	MS	MS	MS
		0-15 cm	30-45 cm	60-75 cm	90-105 cm
Season	1	5286.56*	430.12*	883.19*	16.10
Fence	1	128.17*	1400.72*	3638.91*	5246.06*
Direction	1	0.94	628.81*	1212.39*	1651.31*
Distance	6	6.46	58.14	248.36*	413.71*

\* significant at 0.1%.

**Figure 9.** Effects of snow fences on fall soil moisture at 0-15 cm, at seven distances, leeward and windward of the fence line.




**Figure 10.** Effects of snow fences on fall soil moisture at 30-45 cm, at seven distances, leeward and windward of the fence line.






**Figure 11.** Effects of snow fences on fall soil moisture at 60-75 cm, at seven distances, leeward and windward of the fence line.



**Figure 12.** Effects of snow fences on fall soil moisture at 90-105 cm, at seven distances, leeward and windward of the fence line.



### 3 SPECIES DOCUMENTATION AND SEED BANK

#### 3.1 Species Documentation

A list of the vascular plants of the Oldman River Dam land base is being prepared to determine if there are any rare or endangered species in the land base which must be protected, to identify any species which may be lost due to changes in habitat resulting from the construction of the dam, and to have available for future projects in restoration and revegetation a true reflection of the native vegetation. A preliminary list is included in Appendix II. Voucher specimens were collected and a set will be deposited with the herbaria at the Native Plants Genetic Resources Laboratory (AEC), the Provincial Museum of Alberta (PMAE), and the Oldman River Dam Interpretive Centre. To date, 132 species have been collected (Appendix III). Collection of voucher specimens will continue in 1991 and the Vascular Plants list will be updated.

One hundred and eighty three native plant species have been identified within the land base at the Oldman River Dam. There are another 14 which were reported by Hardy Associates (1986) but which have not yet been collected on-site. These may occur in nearby areas off-site since their study area was larger than the current land base. Voucher specimens were not collected therefore identifications of these species cannot be verified. We have identified 22 species which are introduced. These in most cases are weedy and present some problems for the reestablishment of native plants.

Three species have been found at the Oldman River Dam site which are considered rare in Alberta: *Androsace occidentalis*, *Hymenopappus filifolius* and *Penstemon eriantherus* (Wallis 1987, Wallis et al. 1987, Packer and Bradley 1984). Southwestern Alberta is the northern and western limit of the range over which *A. occidentalis* is found. It is also rare in British Columbia, and to the east in Iowa, Illinois, and endangered in Indiana. This species is however, common in the rest of its range which extends from southern British Columbia, east to western Ontario, and south to California, New Mexico, Texas, and Arkansas (Wallis et al. 1987). On site, *A. occidentalis* is locally scattered under mature Douglas Fir and was found south of Horseshoe Canyon. It will probably survive if left undisturbed although continued heavy grazing in this area may lead to the destruction of this species in the Oldman River Dam land base. Survival could also be in question if the Douglas Fir communities are lost (e.g. due to erosion, etc.).

Both *P. eriantherus* and *H. filifolius* are western species which reach their northern limits in southern Alberta. The former is also rare in British Columbia, Washington, and has a limited western range that includes Idaho, east to North Dakota, Nebraska, and Colorado. *H. filifolius* is native to south Saskatchewan, south to Oregon, Colorado, Kansas, and Mexico (Wallis et al 1987). In the Oldman River Dam land base this species is widely scattered in dry grassland, especially on southern exposures around rocky outcroppings. It was found in an area downstream of the dam and will probably be unaffected by flooding but may be lost due to planned recreational activities in this area. *P. eriantherus* was found only on the south exposures of the hillsides above the work camp. Several plants have already been removed for camp expansion. Since this is the proposed area for a recreational trailer complex, some care must be taken to preserve this small population. Voucher specimens of *A. occidentalis* and *P. eriantherus* have been collected, however none were taken of *H. filifolius* due to the small number of individuals present.

Two other rare species have been reported by Hardy and Associates (1978) Ltd. (1986) on site: *Crataegus douglasii*, and *Osmorhiza chilensis*. The latter is a widespread species which approaches its northwest range limit in Alberta. It is rare in Ontario and South Dakota; uncommon in New England, and threatened in Wisconsin but occurs from southern Alaska, British Columbia, south to California, Arizona, Colorado and east to Newfoundland and south to New England (Wallis et al. 1987).

Alberta represents the northern range of *C. douglasii*. It is a western species with many disjunct populations east (western Ontario, Minnesota, and Michigan) of its range in northern and western North America (Wallis et al 1987). An attempt will be made to document both of these species in 1991 and provide information regarding their preservation in the land base.

### 3.2 Seed Bank

#### 3.2.1 Seed Production

In 1989, the Vegetation Branch (AEC) set up a nursery to multiply seed of some of the native forbs and make them available for various mitigation projects at the Oldman River Dam site. The first crop was harvested in 1990 and will be used in the erosion control plantings being made on the downstream dam face in 1991.

**3.2.2 Seed Collection**

Seed of native grasses, forbs, and shrubs are being collected. These will be maintained at the Native Plant Genetic Resources Laboratory at AEC. Seeds will be multiplied and made available for reclamation projects at the Oldman River Dam site and will be available for future research and selection programs (either at AEC or by other agencies interested in similar goals). A list of the seed collections made in 1990 is included in Appendix IV.



#### 4 REFERENCES CITED

- Aiken, S.G. and S.J. Darbyshire. 1990. Fescue grasses of Canada. Ottawa. 113pp.
- Alberta Environment. 1988a. Oldman River Dam. Wildlife. Oldman River Dam Project Office. Alberta Environment. Edmonton, Alberta. 6pp.
- Alberta Environment. 1988b. Oldman River Dam. Vegetation. Oldman River Dam Project Office. Alberta Environment. Edmonton, Alberta. 6pp.
- Environment Canada. 1989. Daily temperatures and precipitation records for Pincher Creek Airport (April-October, 1989). Unpublished.
- Environment Canada. 1990. Daily temperatures and precipitation records for Pincher Creek Airport. Unpublished.
- Green J.E. and T.R. Eccles. 1989. A strategy for wildlife habitat mitigation for the Oldman Reservoir. The Delta Environmental Management Group. Calgary, Alberta. Prep. for Alberta. 36pp.
- Hardy Associates (1978) Ltd. 1986. Vegetation Inventory: Oldman River Dam. Prepared for Alberta Environment, Planning Division. Calgary, Alberta. 102pp.
- Moss, E.H. 1983. Flora of Alberta. 2nd ed., revised by J.G. Packer. University of Toronto. 687pp.
- Nilson, A.L. and J.E. Green. 1989. An action plan for wildlife habitat mitigation for the Oldman River Dam Reservoir. Prep. for Alberta Public Works Supply and Services. Edmonton, Alberta. 16pp.
- Packer, J.G. and C.E. Bradley. 1984. A checklist of the rare vascular plants in Alberta. Provincial Museum of Alberta. Natural History Occasional Paper No. 5. 112 pp.
- SAS Institute Inc. 1988. SAS/STAT User's Guide: Release 6.03 Edition. 1028 pp.

Smreciu A. 1990. Oldman River Dam, Wildlife Habitat Mitigation. Vegetation Establishment. Interim Progress Report 1989-90. Prep. for Alberta Public Works, Supply and Services. Edmonton, Alberta. 60pp.

Smreciu, E.A. and R.S. Currah. 1989. A guide to the native orchids of Alberta. University of Alberta, Devonian Botanic Garden. 62pp.

Wallis, C. 1987. The rare vascular plants of Alberta: Volume 2. A summary of the taxa occurring in the Canadian Shield, boreal forest, aspen parkland and grassland natural regions. Prep. for Alberta Forestry, Lands and Wildlife. 10pp.

Wallis, C., C. Bradley, M. Fairbarns, and V. Loewen. 1987. The rare flora of Alberta: Volume 3. Species summary sheets. Prep. for Alberta Forestry, Lands and Wildlife.

**Appendix I.** Climate information for Pincher Creek (1989-1990)\*.

Month	Year	Total ppt. (mm)	Max. Temp. (°C)	Min. Temp. (°C)
January	1989	45.6	10.9	-35.0
February	1989	32.4	6.1	-40.1
March	1989	33.0	10.1	-35.1
April	1989	50.2	22.5	-10.5
May	1989	64.8	23.3	-4.1
June	1989	72.8	26.5	2.9
July	1989	73.2	29.8	3.2
August	1989	60.6	30.1	3.5
September	1989	36.8	25.5	-3.7
October	1989	18.2	20.9	-8.3
November	1989	31.3	14.4	-14.5
December	1989	21.7	10.2	-31.0
January	1990	12.4	6.5	-30.1
February	1990	30.6	11.4	-35.0
March	1990	17.6	16.3	-21.9
April	1990	55.0	20.0	-17.9
May	1990	119.8	23.1	-4.2
June	1990	33.8	29.2	0.8
July	1990	76.6	32.6	3.0
August	1990	33.6	32.0	4.4
September	1990	10.2	28.8	-2.2
October	1990	27.4	22.1	-10.7
November	1990	53.6	18.0	-21.2
December	1990	25.2	5.6	-37.1

\* Environment Canada (1989, 1990).

## Appendix II. Vascular Plants of the Oldman River Dam Site.

\* = Introduced.

\*\* = Sightings recorded by Hardy Associates (1978) Ltd. (1986).

### Selaginellaceae

*Selaginella densa* Rydb.

### Club-moss Family

Little Club-moss

### Equisetaceae

*Equisetum arvense* L.

### Horsetail Family

Common Horsetail

### Polypodiaceae

*Cystopteris fragilis* (L.) Bernh.

### Fern Family

Bladder Fern

### Cupressaceae

*Juniperus communis* L.

*Juniperus horizontalis* Moench

### Cypress Family

Ground Juniper

Creeping Juniper

### Pinaceae

*Pinus flexilis* James

*Pseudotsuga menziesii* (Mirb.) Franco

### Pine Family

Limber Pine

Douglas Fir

### Gramineae

*Agropyron dasystachyum* (Hook.) Scribn.

*Agropyron pectiniforme* R. & S.\*

*Agropyron smithii* Rydb.

*Agropyron spicatum* (Pursh) Scribn. & Smith

*Agropyron trachycaulum* (Link) Malte

*Bouteloua gracilis* (HBK) Lag.

*Bromus inermis* Leyss.\*

*Bromus tectorum* L.\*

*Calamagrostis montanensis* Scribn.

*Danthonia parryi* Scribn.

*Elymus piperi* Bowden

*Festuca idahoensis* Elmer\*\*

*Festuca saximontana* Rydb.

*Festuca campestris* Torr.

*Hordeum jubatum* L.\*

*Koeleria macrantha* (Ledeb.) J.A. Schultes f.

*Muhlenbergia cuspidata* (Torr.) Rydb.

*Oryzopsis hymenoides* (R. & S.) Ricker

*Phleum pratense* L.\*

*Poa compressa* L.\*

*Poa palustris* L.

*Poa pratensis* L.

*Stipa comata* Trin. & Rupr.

*Stipa curtisetia* (A.S. Hitchc.) Barkworth

*Stipa viridula* Trin.

### Grass Family

Northern Wheat Grass

Crested Wheat Grass

Western Wheat Grass

Bluebunch Wheat Grass

Slender Wheat Grass

Blue Grama, Buffalo Grass

Awnless Brome

Downy Brome

Plains Reed Grass

Parry Oat Grass

Giant Wild Rye

Bluebunch Fescue

Sheep Fescue

Rough Fescue

Foxtail Barley

June Grass

Muhly Grass

Indian Rice Grass

Timothy

Canada Bluegrass

Fowl Bluegrass

Kentucky Bluegrass

Needle and Thread Grass

Western Porcupine Grass

Green Needle Grass



**Cyperaceae**

*Carex filifolia* Nutt.

*Carex obtusata* Lilj.

**Juncaceae**

*Juncus balticus* Willd.

**Liliaceae**

*Allium cernuum* Roth

*Allium textile* Nels. & Macbr.

*Disporum trachycarpum* (S. Wats.) B. & H. \*\*

*Fritillaria pudica* (Pursh) Spreng.

*Smilacina stellata* (L.) Desf.

*Zigadenus elegans* Pursh

*Zigadenus venenosus* S. Wats

**Iridaceae**

*Sisyrinchium montanum* Greene

**Orchidaceae**

*Coeloglossum viride* (L.) R. Br.

**Salicaceae**

*Populus angustifolia* James

*Populus tremuloides* Michx.

**Betulaceae**

*Alnus* sp.

*Betula occidentalis* Hook.

**Santalaceae**

*Comandra umbellata* (L.) Nutt.

**Polygonaceae**

*Eriogonum flavum* Nutt.

**Nyctaginaceae**

*Mirabilis hirsuta* (Pursh) MacM.

**Caryophyllaceae**

*Cerastium arvense* L.

*Moehringia lateriflora* (L.) Fenzl.

*Paronychia sessiliflora* Nutt.

*Silene noctiflora* L.\*

**Ranunculaceae**

*Anemone cylindrica* A. Gray

*Anemone multifida* Poir.

**Sedge Family**

Thread-leaved Sedge

Blunt Sedge

**Rush Family**

Wire Rush

**Lily Family**

Nodding Onion

Prairie Onion

Fairy-bells

Yellow-bell

Star-flowered Solomon's-seal

White Camas

Death Camas

**Iris Family**

Blue-eyed Grass

**Orchid Family**

Bracted Orchid

**Willow Family**

Narrow-leaf Cottonwood

Aspen

**Birch Family**

Alder

Water Birch; Black Birch

**Sandalwood Family**

Bastard Toadflax

**Buckwheat Family**

Yellow Umbrella-plant

**Four-o'clock Family**

Umbrellawort

**Pink Family**

Mouse-ear Chickweed

Sandwort

Low Whitlow-wort

Night-flowering Catchfly

**Crowfoot Family**

Long-fruited Anemone

Cut-leaved Anemone

*Anemone patens* L.  
*Clematis ligusticifolia* Nutt.  
*Delphinium bicolor* Nutt.  
*Thalictrum venulosum* Trel.

#### Cruciferae

*Arabis divaricarpa* L.  
*Arabis nuttallii* Robinson  
*Conringia orientalis* (L.) Dum.\*  
*Erysimum cheiranthoides* L.\*  
*Erysimum inconspicuum* (S. Wats.) MacM.  
*Lesquerella arenosa* (Richards.) Rydb.  
*Physaria didymocarpa* (Hook.) A. Gray  
*Thlaspi arvense* L.\*,\*\*

#### Saxifragaceae

*Heuchera parvifolia* Nutt. ex T. & G.

#### Grossulariaceae

*Ribes hirtellum* Michx.\*\*  
*Ribes oxycanthoides* L.

#### Rosaceae

*Amelanchier alnifolia* Nutt.  
*Chamaerhodos erecta* (L.) Bunge  
*Crataegus douglasii* Lindl.  
*Crataegus rotundifolia* Moench  
*Fragaria virginiana* Duchesne  
*Geum allepicum* Jacq.  
*Geum triflorum* Pursh  
*Potentilla anserina* L.  
*Potentilla concinna* Richards.  
*Potentilla fruticosa* L.  
*Potentilla gracilis* Dougl. ex Hook.  
*Potentilla hippiana* Lehm.  
*Potentilla pensylvanica* L.  
*Prunus virginiana* L.  
*Rosa acicularis* Lindl.  
*Rosa arkansana* Porter  
*Rosa woodsii* Lindl.

#### Leguminosae

*Astragalus bisulcatus* (Hook.) A. Gray  
*Astragalus crassicaarpus* Nutt.  
*Astragalus dasyglottis* Fisch. ex DC.  
*Astragalus drummondii* Dougl. ex Hook.  
*Astragalus flexuosus* Dougl.  
*Astragalus gilviflorus* Sheldon

Prairie Crocus  
Western Clematis  
Low Larkspur  
Veiny Meadow-rue

#### Mustard Family

Rock Cress  
Rock Cress  
Hare's-ear Mustard  
Wormseed Mustard  
Small-flowered Rocket  
Bladder-pod  
Double Bladder-pod  
Pennycress; Stinkweed

#### Saxifrage Family

Alum-root

#### Gooseberry Family

Wild Gooseberry  
Wild Gooseberry

#### Rose Family

Saskatoon  
Chamaerhodos  
Douglas Hawthorn  
Round-leaf Hawthorn  
Wild Strawberry  
Yellow Avens  
Three-flowered Avens  
Silverweed  
Early Cinquefoil  
Shrubby Cinquefoil  
Graceful Potentilla  
Wooly Cinquefoil  
Prairie Cinquefoil  
Choke Cherry  
Prickly Rose  
Prairie Rose  
Common Wild Rose

#### Pea Family

Two-grooved Milk Vetch  
Ground Plum  
  
Drummond's Milk Vetch  
Slender Milk Vetch  
Cushion Milk Vetch

*Astragalus missouriensis* Nutt.  
*Astragalus pectinatus* Dougl. ex Hook.  
*Astragalus tenellus* Pursh  
*Astragalus vexilliflexus* Sheldon  
*Glycyrrhiza lepidota* (Nutt.) Pursh  
*Hedysarum alpinum* L.  
*Hedysarum boreale* Nutt.  
*Hedysarum sulphurescens* Rydb.  
*Lupinus argenteus* Pursh  
*Lupinus sericeus* Pursh  
*Medicago sativa* L.\*  
*Melilotus alba* Desr.\*  
*Melilotus officinalis* (L.) Lam.\*  
*Oxytropis monticola* A. Gray  
*Oxytropis sericea* Nutt.  
*Oxytropis splendens* Dougl. ex Hook.  
*Petalostemon candidum* (Willd.) Michx.  
*Petalostemon purpureum* (Vent.) Rydb.  
*Psoralea esculenta* Pursh  
*Thermopsis rhombifolia* (Nutt.) Richards.  
*Vicia americana* Muhl.

**Geraniaceae**

*Geranium viscosissimum* Fisch. & Mey.

**Linaceae**

*Linum lewisii* Pursh  
*Linum rigidum* Pursh

**Anacardiaceae**

*Rhus trilobata* Nutt.

**Malvaceae**

*Sphaeralcea coccinea* (Pursh) Rydb.

**Violaceae**

*Viola adunca* J.E. Smith  
*Viola canadensis* L.  
*Viola nuttallii* Pursh  
*Viola renifolia* A. Gray\*\*

**Loasaceae**

*Mentzelia decapetala* (Sims) Urban & Gilg

**Cactaceae**

*Coryphantha vivipara* (Nutt.) Britt. & Rose  
*Opuntia polyacantha* Haw.

**Elaeagnaceae**

Missouri Milk Vetch  
Narrow-leaved Milk Vetch  
Loose-flowered Milk Vetch  
Bent-flowered Milk Vetch  
Wild Licorice  
American Sweetbroom  
Northern Hedysarum  
Yellow Hedysarum  
Perennial Lupine  
Perennial Lupine  
Alfalfa  
White Sweet Clover  
Yellow Sweet Clover  
Late Yellow Loco-weed  
Early Yellow Loco-weed  
Showy Locoweed  
White Prairie Clover  
Purple Prairie Clover  
Indian Bread-root  
Buffalo Bean  
Wild Vetch

**Geranium Family**

Sticky Purple Geranium

**Flax Family**

Wild Blue Flax  
Yellow Flax

**Sumach Family**

Skunk-bush

**Mallow Family**

Scarlet Mallow

**Violet Family**

Early Blue Violet  
Western Canada Violet  
Yellow Prairie Violet  
Kidney-leaved Violet

**Loasa Family**

Evening Star

**Cactus Family**

Cushion; Ball Cactus  
Prickly Pear

**Oleaster Family**



*Elaeagnus commutata* Bernh. ex Rydb.  
*Shepherdia argentea* Nutt.\*\*  
*Shepherdia canadensis* (L.) Nutt.

#### Onagraceae

*Gaura coccinea* Pursh  
*Oenothera biennis* L.  
*Oenothera nuttallii* Sweet\*\*

#### Umbelliferae

*Heracleum lanatum* Michx.\*\*  
*Lomatium triternatum* (Pursh) Coult. & Rose  
*Musineon divaricatum* (Pursh) Nutt.  
*Zizia aptera* (A. Gray) Fern.

*Osmorhiza chilensis* Hook. & Arn.\*\*  
*Perideridia gairdneri* (Hook. & Arn.) Mathius

#### Cornaceae

*Cornus stolonifera* Michx.

#### Ericaceae

*Arctostaphylos uva-ursi* (L.) Spreng

#### Primulaceae

*Androsace chamaejasme* Host  
*Androsace occidentalis* Pursh  
*Androsace septentrionalis* L.  
*Dodecatheon conjugens* Greene

#### Gentianaceae

*Gentianella amarella* (L.) Borner

#### Apocynaceae

*Apocynum androsaemifolium* L.  
*Apocynum cannabinum* L.\*\*

#### Polemoniaceae

*Collomia linearis* Nutt.  
*Phlox hoodii* Richards  
*Polemonium pulcherrimum* Hook.

#### Boraginaceae

*Cryptantha nubigena* (Greene) Payson  
*Cynoglossum officinale* L.\*  
*Hackelia floribunda* (Lehm.) I.M. Johnston  
*Lappula occidentalis* (S. Wats.) Greene\*  
*Lappula squarrosa* Moench\*

Wolf Willow  
Thorny Buffalo-berry  
Canadian Buffalo-berry

#### Evening Primrose Family

Scarlet Butterfly-weed  
Yellow Evening Primrose  
White Evening Primrose

#### Carrot Family

Cow Parsnip  
Prairie Parsley  
Leafy Musineon  
Meadow Parsnip, Heartleafed  
Alexanders  
Sweet Cicely  
Squaw Root, Yampa

#### Dogwood Family

Red-osier Dogwood

#### Heath Family

Common Bearberry,  
Kinnickinnick

#### Primrose Family

Sweet-flowered Androsace  
Fairy Candelabra  
Fairy Candelabra  
Shooting Star

#### Gentian Family

Felwort

#### Dogbane Family

Spreading Dogbane  
Indian Hemp

#### Phlox Family

Collomia  
Moss Phlox  
Jacobs-ladder

#### Borage Family

CLustered Oreocarya  
Hound's-tongue  
Stick-seed  
Stick-tights  
Beggar-ticks



*Lithospermum incisum* Lehm.  
*Lithospermum ruderae* Lehm.

**Labiatae**

*Mentha arvensis* L.  
*Monarda fistulosa* L.  
*Stachys palustris* L.\*\*

**Scrophulariaceae**

*Besseyia wyomingensis* (A. Nels) Rydb.  
*Castilleja lutescens* (Greenm.) Rydb.  
*Orthocarpus luteus* Nutt.  
*Penstemon confertus* Dougl.  
*Penstemon eriantherus* Pursh  
*Penstemon nitidus* Dougl. ex Benth.  
*Penstemon procerus* Dougl. ex Grah.

**Orobanchaceae**

*Orobanche fasciculata* Nutt.

**Plantaginaceae**

*Plantago canescens* Adams

**Rubiaceae**

*Galium boreale* L.

**Caprifoliaceae**

*Symphoricarpos albus* (L.) Blake\*\*  
*Symphoricarpos occidentalis* Hook.

**Campanulaceae**

*Campanula rotundifolia* L.

**Compositae**

*Achillea millefolium* L.  
*Agoseris glauca* (Pursh) Raf  
*Antennaria anaphaloides* Rydb.  
*Antennaria parvifolia* Nutt.

*Antennaria rosea* Greene  
*Arnica fulgens* Pursh\*\*  
*Arnica sororia* Greene  
*Artemisia campestris* L.  
*Artemisia dracunculoides* L.\*\*  
*Artemisia frigida* Willd.  
*Artemisia ludoviciana* Nutt.  
*Aster conspicuus* Lindl.\*\*  
*Aster ericoides* L.  
*Aster falcatus* Lindl.

Puccoon  
Yellow Puccoon

**Mint Family**

Wild Mint  
Wild Bergamot  
Hedge Nettle

**Figwort Family**

Kitten-tails  
Yellow Indian Paint-brush  
Owl-clover  
Yellow Beard-tongue  
Crested Beard-tongue  
Smooth Blue Beard-tongue  
Slender Blue Beard-tongue

**Broom-rape Family**

Clustered Broom-rape

**Plantain Family**

Plantain

**Madder Family**

Northern Bedstraw

**Honeysuckle Family**

Snowberry  
Buckbrush

**Bluebell Family**

Bluebell; Harebell

**Composite Family**

Common Yarrow  
False Dandelion  
Pussy Toes, Tall Everlasting  
Pussy Toes, Small-leaved  
Everlasting  
Pink Pussy Toes  
Shining Arnica  
Twin Arnica  
Field Sage  
Dragonwort  
Pasture Sage  
Prairie Sage  
Showy Aster  
Tufted White Prairie Aster  
Creeping White Prairie Aster

<i>Aster laevis</i> L.	Smooth Aster
<i>Balsamorhiza sagittata</i> (Pursh) Nutt.	Balsam-root
<i>Cirsium arvense</i> (L.) Scop.*	Canada Thistle
<i>Cirsium undulatum</i> (Nutt.) Spreng.	Wavy-leaved Thistle
<i>Erigeron caespitosus</i> Nutt.	Tufted Fleabane
<i>Erigeron compositus</i> Pursh	Compound Fleabane
<i>Erigeron glabellus</i> Nutt.	Smooth Fleabane
<i>Erigeron speciosus</i> (Lindl.) DC.	Showy Fleabane
<i>Gaillardia aristata</i> Pursh	Gaillardia; Blanket Flower
<i>Grindelia squarrosa</i> (Pursh) Dunal	Gumweed
<i>Gutierrezia sarothrae</i> (Pursh) Britt. & Rusby	Broomweed
<i>Haplopappus lanceolatus</i> (Hook.) T. & G.**	Lance-leaved Pyrocoma
<i>Haplopappus spinulosus</i> (Pursh) DC.	Spiny Ironplant
<i>Helianthus annuus</i> L.	Common Annual Sunflower
<i>Helianthus subrhomboides</i> Rydb.	Rhombic-leaved Sunflower
<i>Heterotheca villosa</i> (Pursh) Shinnery	Golden Aster
<i>Hymenopappus filifolius</i> Hook.	Tufted Hymenopappus
<i>Hymenoxys acaulis</i> (Pursh) Parker	Stemless Rubberweed
<i>Hymenoxys richardsonii</i> (Hook.) Cockerell	Colorado Rubber-plant
<i>Liatris punctata</i> Hook.	Dotted Blazing Star
<i>Lygodesmia juncea</i> (Pursh) D. Don	Skeleton-weed
<i>Ratibida columnifera</i> (Nutt.) Wooten & Standl.	Prairie Cone-flower
<i>Senecio canus</i> Hook.	Prairie Groundsel
<i>Senecio integerrimus</i> Nutt.	Entire-leaved Groundsel
<i>Solidago missouriensis</i> Nutt.	Missouri Goldenrod
<i>Solidago rigida</i> L.	Stiff Goldenrod
<i>Taraxacum officinale</i> Weber*	Dandelion
<i>Townsendia exscapa</i> (Richards.) Porter	Townsendia
<i>Tragopogon dubius</i> Scop.*	Goat's-beard
<i>Xanthium strumarium</i> L.*	Cocklebur

**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
AS098	<i>Achillea millefolium</i>	07-06-90	S19-R29-Twp7-W4
AS141	<i>Achillea millefolium</i>	18-06-90	S11-R1-Twp7-W5
JH158	<i>Agoseris glauca</i>	26-06-90	S16-R29-Twp7-W4
AS136	<i>Agropyron spicatum</i>	14-06-90	S32-R1-Twp7-W5
AS199	<i>Agropyron dasystachyum</i>	09-08-90	S32-R1-Twp7-W4
AS111	<i>Agropyron smithii</i>	08-06-90	S16-R29-Twp7-W4
AS137	<i>Agropyron smithii</i>	14-06-90	S32-R1-Twp7-W5
AS191	<i>Agropyron smithii</i>	09-08-90	S32-R1-Twp7-W4
AS011	<i>Agropyron spicatum</i>	18-07-89	S32-R1-Twp7-W5
AS193	<i>Agropyron spicatum</i>	09-08-90	S32-R1-Twp7-W4
AS135	<i>Agropyron trachycaulum?</i>	14-06-90	S32-R1-Twp7-W5
AS198	<i>Agropyron trachycaulum?</i>	09-08-90	S32-R1-Twp7-W4
AS003	<i>Allium cernuum</i>	18-07-89	S32-R1-Twp7-W5
JH181	<i>Allium cernuum</i>	20-07-90	S19-R29-Twp7-W4
AS184	<i>Allium cernuum</i>	09-08-90	S32-R1-Twp7-W4
AS071	<i>Allium textile</i>	28-06-90	S16-R29-Twp7-W4
AS090	<i>Androsace occidentalis</i>	07-06-90	S24-R1-Twp7-W5
AS065	<i>Anemone multifida</i>	28-06-90	S16-R29-Twp7-W4
AS094	<i>Antennaria anaphaloides</i>	07-06-90	S19-R29-Twp7-W4
AS054	<i>Antennaria parvifolia</i>	28-06-90	S11-R1-Twp7-W5
AS014	<i>Antennaria parvifolia</i>	18-07-89	S32-R1-Twp7-W5
AS049	<i>Antennaria rosea</i>	28-06-90	S11-R1-Twp7-W5
JH178	<i>Apocynum androsaemifolium</i>	19-07-90	S16-R29-Twp7-W4
AS082	<i>Arabis divaricarpa</i>	06-06-90	S32-R1-Twp7-W5
AS100	<i>Arabis divaricarpa</i>	07-06-90	S19-R29-Twp7-W4
AS038	<i>Arabis nuttallii</i>	06-06-90	S32-R1-Twp7-W5
AS081	<i>Arabis nuttallii</i>	06-06-90	S32-R1-Twp7-W5

**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
AS099	<i>Arabis nuttallii</i>	07-06-90	S19-R29-Twp7-W4
AS055	<i>Arabis nuttallii</i>	28-06-90	S11-R1-Twp7-W5
AS031	<i>Arabis nuttallii</i>	06-06-90	S32-R1-Twp7-W5
AS140	<i>Arnica sororia</i>	14-06-90	S32-R1-Twp7-W5
AS188	<i>Artemisia campestris</i>	09-08-90	S32-R1-Twp7-W4
JH213	<i>Artemisia campestris</i>	27-08-90	S17-R29-Twp7-W4
AS215	<i>Aster ericoides</i>	27-08-90	S17-R29-Twp7-W4
AS186	<i>Aster laevis</i>	09-08-90	S32-R1-Twp7-W4
JH214	<i>Aster laevis</i>	27-08-90	S17-R29-Twp7-W4
AS142	<i>Astragalus bisulcatus</i>	18-06-90	S11-R1-Twp7-W5
AS068	<i>Astragalus crassicaarpus</i>	28-06-90	S16-R29-Twp7-W4
AS109	<i>Astragalus drummondii</i>	08-06-90	S16-R29-Twp7-W4
AS119	<i>Astragalus flexuosus</i>	12-06-90	S17-R29-Twp7-W4
AS020	<i>Astragalus gilviflorus</i>	06-06-90	S33-R1-Twp7-W5
AS073	<i>Astragalus missouriensis</i>	28-06-90	S16-R29-Twp7-W4
AS021	<i>Astragalus missouriensis</i>	06-06-90	S16-R29-Twp7-W4
AS106	<i>Astragalus pectinatus</i>	08-06-90	S16-R29-Twp7-W4
AS107	<i>Astragalus striatus?</i>	08-06-90	S16-R29-Twp7-W4
AS168	<i>Astragalus striatus?</i>	28-06-90	S16-R29-Twp7-W4
AS102	<i>Astragalus tenellus</i>	07-06-90	S19-R29-Twp7-W4
AS057	<i>Astragalus tenellus</i>	28-06-90	S16-R29-Twp7-W4
AS077	<i>Astragalus vexilliflexus</i>	06-06-90	S24-R1-Twp7-W5
AS074	<i>Astragalus vexilliflexus</i>	28-06-90	S16-R29-Twp7-W4
AS030	<i>Balsamorhiza sagittata</i>	06-06-90	S16-R29-Twp7-W4
AS029	<i>Besseyia wyomingensis</i>	06-06-90	S33-R1-Twp7-W5
AS138	<i>Besseyia wyomingensis</i>	14-06-90	S32-R1-Twp7-W5
AS133	<i>Betula occidentalis</i>	14-06-90	S16-R29-Twp7-W4



**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
AS203	<i>Bouteloua gracilis</i>	10-08-90	S16-R29-Twp7-W4
AS072	<i>Calamagrostis montanensis?</i>	28-06-90	S16-R29-Twp7-W4
JH159	<i>Campanula rotundifolia</i>	26-06-90	S16-R29-Twp7-W4
AS078	<i>Carex filifolia</i>	06-06-90	S24-R1-Twp7-W5
AS032	<i>Carex obtusata</i>	06-06-90	S33-R1-Twp7-W5
AS008	<i>Carex</i> sp.	18-07-89	S32-R1-Twp7-W5
AS009	<i>Carex</i> sp.	18-07-89	S32-R1-Twp7-W5
AS096	<i>Castilleja lutescens</i>	07-06-90	S19-R29-Twp7-W4
AS037	<i>Cerastium arvense</i>	06-06-90	S33-R1-Twp7-W5
AS041	<i>Cerastium arvense</i>	28-06-90	S11-R1-Twp7-W5
AS087	<i>Chamaerhodos erecta</i>	07-06-90	S19-R29-Twp7-W4
JH179	<i>Clematis ligusticifolia</i>	19-07-90	S16-R29-Twp7-W4
AS122	<i>Coeloglossum viride</i>	14-06-90	S16-R1-Twp7-W4
AS043	<i>Comandra umbellata</i>	28-06-90	S11-R1-Twp7-W5
AS002	<i>Conringia orientalis</i>	29-06-89	S32-R1-Twp7-W4
AS165	<i>Cornus stolonifera</i>	28-06-90	S16-R29-Twp7-W4
AS105	<i>Crataegus rotundifolia</i>	08-06-90	S16-R29-Twp7-W4
AS110	<i>Crucifer</i> (weed)	08-06-90	S16-R29-Twp7-W4
AS085	<i>Cryptantha nubigena</i>	07-06-90	S19-R29-Twp7-W4
AS069	<i>Cryptantha nubigena</i>	28-06-90	S16-R29-Twp7-W4
AS134	<i>Cystopteris fragilis</i>	14-06-90	S16-R29-Twp7-W4
AS125	<i>Danthonia parryi</i>	14-06-90	S16-R29-Twp7-W4
AS192	<i>Danthonia parryi</i>	09-08-90	S32-R1-Twp7-W4
AS023	<i>Delphinium bicolor</i>	06-06-90	S16-R29-Twp7-W4
AS060	<i>Delphinium bicolor</i>	28-06-90	S16-R29-Twp7-W4
AS025	<i>Dodecatheon conjugens</i>	06-06-90	S32-R1-Twp7-W5

**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
AS123	<i>Elaeagnus commutata</i>	14-06-90	S16-R29-Twp7-W4
AS004	<i>Elymus piperi</i>	18-07-89	S16-R29-Twp7-W4
JH175	<i>Erigeron caespitosus</i>	19-07-90	S16-R29-Twp7-W4
AS190	<i>Erigeron caespitosus</i>	09-08-90	S32-R1-Twp7-W4
JH182	<i>Erigeron speciosus</i>	20-07-90	S19-R29-Twp7-W4
AS152	<i>Eriogonum flavum</i>	21-06-90	S17-R29-Twp7-W4
AS084	<i>Erysimum cheiranthoides</i>	07-06-90	S19-R29-Twp7-W4
AS115	<i>Festuca saximontana?</i>	08-06-90	S16-R29-Twp7-W4
AS195	<i>Festuca saximontana?</i>	09-08-90	S32-R1-Twp7-W4
AS101	<i>Festuca campestris ?</i>	07-05-90	S19-R29-Twp7-W4
AS080	<i>Festuca campestris</i>	06-06-90	S24-R1-Twp7-W5
AS092	<i>Festuca campestris</i>	07-06-90	S19-R29-Twp7-W4
AS114	<i>Festuca campestris</i>	08-06-90	S16-R29-Twp7-W4
AS075	<i>Festuca campestris</i>	28-06-90	S16-R29-Twp7-W4
AS056	<i>Festuca campestris</i>	28-06-90	S16-R29-Twp7-W4
AS196	<i>Festuca campestris</i>	09-08-90	S32-R1-Twp7-W4
AS034	<i>Festuca campestris?</i>	06-06-90	S32-R1-Twp7-W5
AS036	<i>Festuca campestris?</i>	06-06-90	S16-R29-Twp7-W4
AS051	<i>Fragaria virginiana</i>	28-06-90	S11-R1-Twp7-W5
AS026	<i>Fritillaria pudica</i>	06-06-90	S16-R29-Twp7-W4
AS150	<i>Gaillardia aristata</i>	21-06-90	S17-R29-Twp7-W4
JH177	<i>Galium boreale</i>	19-07-90	S16-R29-Twp7-W4
AS162	<i>Gaura coccinea</i>	27-06-90	S19-R29-Twp7-W4
AS167	<i>Gaura coccinea</i>	28-06-90	S16-R29-Twp7-W4
AS209	<i>Gentianella amarella</i>	10-08-90	S19-R29-Twp7-W4
AS121	<i>Geranium viscosissimum</i>	12-06-90	S17-R29-Twp7-W4
AS027	<i>Geum triflorum</i>	06-06-90	S33-R1-Twp7-W5

**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
AS045	<i>Geum triflorum</i>	28-06-90	S11-R1-Twp7-W5
AS128	Grass sp.	14-06-90	S16-R29-Twp7-W4
AS211	<i>Poa palustris</i>	10-08-90	S19-R29-Twp7-W4
AS205	<i>Grindelia squarrosa</i>	10-08-90	S16-R29-Twp7-W4
AS216	<i>Gutierrezia sarothrae</i>	27-08-90	S17-R29-Twp7-W4
AS079	<i>Hackelia floribunda</i>	06-06-90	S24-R1-Twp7-W5
AS089	<i>Hackelia floribunda</i>	07-06-90	S19-R29-Twp7-W4
JH212	<i>Haplopappus spinulosus</i>	27-08-90	S17-R29-Twp7-W4
AS130	<i>Hedysarum alpinum</i>	14-06-90	S16-R29-Twp7-W4
AS129	<i>Hedysarum sulphurescens</i>	14-06-90	S16-R29-Twp7-W4
AS067	<i>Hedysarum sulphurescens</i>	28-06-90	S16-R29-Twp7-W4
JH183	<i>Helianthus annuus</i>	20-07-90	S17-R29-Twp7-W4
JH174	<i>Helianthus subrhomboides</i>	19-07-90	S16-R29-Twp7-W4
JH172	<i>Heterotheca villosa</i>	19-07-90	S16-R29-Twp7-W4
AS118	<i>Heuchera parvifolia</i>	08-06-90	S27-R30-Twp7-W4
AS066	<i>Hymenoxys acaulis</i>	28-06-90	S16-R29-Twp7-W4
JH157	<i>Hymenoxys richarsonii</i>	26-06-90	S16-R29-Twp7-W4
AS153	<i>Koeleria macrantha</i>	21-06-90	S17-R29-Twp7-W4
AS035	<i>Koeleria macrantha?</i>	06-06-90	S16-R29-Twp7-W4
AS093	<i>Koeleria macrantha?</i>	07-06-90	S19-R29-Twp7-W4
AS112	<i>Koeleria macrantha</i>	08-06-90	S16-R29-Twp7-W4
AS048	<i>Lesquerella arenosa</i>	28-06-90	S11-R1-Twp7-W5
AS197	<i>Liatris punctata</i>	09-08-90	S32-R1-Twp7-W4
AS217	<i>Liatris punctata</i>	27-08-90	S17-R29-Twp7-W4
AS086	<i>Linum lewisii</i>	07-06-90	S19-R29-Twp7-W4
AS143	<i>Linum lewisii</i>	21-06-90	S17-R29-Twp7-W4
JH155	<i>Linum rigidum</i>	26-06-90	S16-R29-Twp7-W4



**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
AS058	<i>Lithospermum incisum</i>	28-06-90	S16-R29-Twp7-W4
AS046	<i>Lithospermum ruderales</i>	28-06-90	S11-R1-Twp7-W5
AS033	<i>Lomatium triternatum</i>	06-06-90	S33-R1-Twp7-W5
AS059	<i>Lomatium triternatum</i>	28-06-90	S16-R29-Twp7-W4
AS063	<i>Lupinus argenteus</i>	28-05-90	S16-R29-Twp7-W4
AS116	<i>Lupinus argenteus</i>	08-06-90	S16-R29-Twp7-W4
AS117	<i>Lupinus sericeus?</i>	08-06-90	S16-R29-Twp7-W4
AS207	<i>Lygodesmia juncea</i>	10-08-90	S16-R29-Twp7-W4
AS204	<i>Menztelesia decapetala</i>	10-08-90	S16-R29-Twp7-W4
AS132	<i>Moehringia lateriflora</i>	14-06-90	S16-R29-Twp7-W4
JH170	<i>Monarda fistulosa</i>	19-07-90	S16-R29-Twp7-W4
AS202	<i>Muhlenbergia cuspidata</i>	09-08-90	S16-R29-Twp7-W4
AS019	<i>Musineon divaricatum</i>	06-06-90	S33-R1-Twp7-W5
AS050	<i>Musineon divaricatum</i>	28-06-90	S11-R1-Twp7-W5
AS145	<i>Orobancha fasciculata</i>	21-06-90	S17-R29-Twp7-W4
JH176	<i>Orthocarpus luteus</i>	19-07-90	S16-R29-Twp7-W4
AS187	<i>Orthocarpus luteus</i>	09-08-90	S32-R1-Twp7-W4
AS163	<i>Oryzopsis hymenoides</i>	27-06-90	S19-R29-Twp7-W4
AS091	<i>Oxytropis sericea</i>	07-06-90	S19-R29-Twp7-W4
AS053	<i>Oxytropis sericea</i>	28-06-90	S11-R1-Twp7-W5
AS146	<i>Paronychia sessiliflora</i>	21-06-90	S17-R29-Twp7-W4
AS139	<i>Penstemon confertus</i>	14-06-90	S32-R1-Twp7-W5
AS001	<i>Penstemon eriantherus</i>	14-06-89	S17-R29-Twp7-W4
AS120	<i>Penstemon eriantherus</i>	12-06-90	S17-R29-Twp7-W4
AS088	<i>Penstemon nitidus</i>	07-06-90	S19-R29-Twp7-W4
AS047	<i>Penstemon nitidus</i>	28-06-90	S11-R1-Twp7-W5
AS208	<i>Perideridia gairdneri</i>	10-08-90	S19-R29-Twp7-W4



**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
JH169	<i>Petalostemon candidum</i>	19-07-90	S16-R29-Twp7-W4
JH173	<i>Petalostemon purpureum</i>	19-07-90	S16-R29-Twp7-W4
AS185	<i>Petalostemon purpureum</i>	09-08-90	S32-R1-Twp7-W4
AS028	<i>Phlox hoodii</i>	06-06-90	S33-R1-Twp7-W5
AS024	<i>Physaria didymocarpa</i>	06-06-90	S16-R29-Twp7-W4
AS070	<i>Physaria didymocarpa</i>	28-06-90	S16-R29-Twp7-W4
AS131	<i>Plantago canescens</i>	14-06-90	S16-R29-Twp7-W4
AS200	<i>Poa compressa</i>	09-08-90	S16-R29-Twp7-W4
AS005	<i>Poa pratensis</i>	18-07-89	S16-R29-Twp7-W4
AS042	<i>Poa pratensis?</i>	28-06-90	S11-R1-Twp7-W5
AS052	<i>Poa pratensis?</i>	28-06-90	S11-R1-Twp7-W5
AS194	<i>Poa pratensis?</i>	09-08-90	S32-R1-Twp7-W4
AS113	<i>Poa</i> sp.	08-06-90	S16-R29-Twp7-W4
AS076	<i>Polemonium pulcherrimum</i>	06-06-90	S24-R1-Twp7-W5
AS164	<i>Populus angusticifolia</i>	28-06-90	S16-R29-Twp7-W4
AS040	<i>Potentilla concinna</i>	06-06-90	S33-R1-Twp7-W5
JH156	<i>Potentilla fruticosa</i>	26-06-90	S16-R29-Twp7-W4
AS149	<i>Potentilla hippiana</i>	21-06-90	S17-R29-Twp7-W4
AS013	<i>Potentilla hippiana</i>	18-07-89	S16-R29-Twp7-W4
AS151	<i>Potentilla pensylvanica</i>	21-06-90	S17-R29-Twp7-W4
AS161	<i>Potentilla pensylvanica</i>	27-06-90	S19-R29-Twp7-W4
AS012	<i>Potentilla pensylvanica</i>	18-07-89	S16-R29-Twp7-W4
AS104	<i>Prunus virginiana</i>	08-06-90	S16-R29-Twp7-W4
AS147	<i>Psoralea esculenta</i>	21-06-90	S17-R29-Twp7-W4
JH180	<i>Ratibida columnifera</i>	20-07-90	S19-R29-Twp7-W4
AS062	<i>Rhus trilobata</i>	28-06-90	S16-R29-Twp7-W4
AS126	<i>Rosa arkansana</i>	14-06-90	S16-R29-Twp7-W4

**Appendix III.** Voucher specimens collected (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
AS166	<i>Rosa arkansana</i>	28-06-90	S33-R1-Twp7-W5
AS097	<i>Senecio canus</i>	07-06-90	S19-R29-Twp7-W4
AS064	<i>Senecio canus</i>	28-06-90	S16-R29-Twp7-W4
AS083	<i>Senecio integerrimus</i>	07-06-90	S19-R29-Twp7-W4
AS044	<i>Shepherdia canadensis</i>	28-06-90	S11-R1-Twp7-W5
JH160	<i>Sisyrinchium montanum</i>	26-06-90	S16-R29-Twp7-W4
AS061	<i>Smilacina stellata</i>	28-06-90	S16-R29-Twp7-W4
AS006	<i>Solidago missouriensis</i>	13-07-89	S11-R1-Twp7-W5
AS007	<i>Solidago missouriensis</i>	18-07-89	S16-R29-Twp7-W5
JH171	<i>Solidago missouriensis</i>	19-07-90	S16-R29-Twp7-W4
AS189	<i>Solidago missouriensis</i>	09-08-90	S32-R1-Twp7-W4
AS206	<i>Solidago missouriensis</i>	10-08-90	S16-R29-Twp7-W4
AS144	<i>Sphaeralcea coccinea</i>	21-06-90	S17-R29-Twp7-W4
AS154	<i>Stipa comata</i>	21-06-90	S17-R29-Twp7-W4
AS010	<i>Stipa comata</i>	18-07-89	S32-R1-Twp7-W5
AS201	<i>Stipa comata</i>	09-08-90	S16-R29-Twp7-W4
AS210	<i>Symphoricarpos occidentalis</i>	10-08-90	S19-R29-Twp7-W4
AS039	<i>Thermopsis rhombifolia</i>	06-06-90	S33-R1-Twp7-W5
AS148	<i>Vicia americana</i>	21-06-90	S17-R29-Twp7-W4
AS103	<i>Viola adunca</i>	07-06-90	S19-R29-Twp7-W4
AS218	<i>Viola canadensis</i>	28-05-90	S16-R29-Twp7-W4
AS022	<i>Viola nuttallii</i>	06-06-90	S16-R29-Twp7-W4
AS108	<i>Viola nuttallii</i>	08-06-90	S16-R29-Twp7-W4
AS127	<i>Zigadenus elegans</i>	14-06-90	S16-R29-Twp7-W4
AS095	<i>Zigadenus venenosus</i>	07-06-90	S19-R29-Twp7-W4
AS124	<i>Zizia aptera</i>	14-06-90	S16-R29-Twp7-W4

**Appendix IV.** Native seed collection (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
88	<i>Agoseris glauca</i>	06-06-90	S19-R29-Twp7-W4
100	<i>Agropyron smithii</i>	09-08-90	S32-R1-Twp7-W5
98	<i>Agropyron spicatum</i>	09-08-90	S32-R1-Twp7-W5
140	<i>Allium cernuum</i>	27-08-90	S16-R29-Twp7-W4
79	<i>Allium textile</i>	19-07-90	S16-R29-Twp7-W4
65	<i>Allium textile</i>	21-07-90	S17-R29-Twp7-W4
89	<i>Anemone multifida</i>	19-07-90	S16-R29-Twp7-W4
164	<i>Anemone patens</i>	07-06-90	S19-R29-Twp7-W4
47	<i>Antennaria parvifolia</i>	27-06-90	S17-R29-Twp7-W4
150	<i>Aster laevis</i>	27-08-90	S16-R29-Twp7-W4
60	<i>Astragalus drummondii</i>	21-07-90	S16-R29-Twp7-W4
132	<i>Astragalus drummondii</i>	10-08-90	S16-R29-Twp7-W4
62	<i>Astragalus flexuosus</i>	21-07-90	S17-R29-Twp7-W4
122	<i>Astragalus flexuosus</i>	09-08-90	S32-R1-Twp7-W5
149	<i>Astragalus gilvaeflorus</i>	27-08-90	S17-R29-Twp7-W4
86	<i>Astragalus missouriensis</i>	10-07-90	S16-R29-Twp7-W4
76	<i>Astragalus missouriensis</i>	19-07-90	S16-R29-Twp7-W4
61	<i>Astragalus missouriensis</i>	21-07-90	S16-R29-Twp7-W4
67	<i>Astragalus missouriensis</i>	21-07-90	S17-R29-Twp7-W4
131	<i>Astragalus missouriensis</i>	10-08-90	S16-R29-Twp7-W4
118	<i>Astragalus pectinatus</i>	08-08-90	S19-R29-Twp7-W4
120	<i>Astragalus striatus</i>	08-08-90	S19-R29-Twp7-W4
128	<i>Astragalus striatus</i>	09-08-90	S17-R29-Twp7-W4
153	<i>Astragalus striatus</i>	27-08-90	S16-R29-Twp7-W4
87	<i>Astragalus tenellus</i>	19-07-90	S16-R29-Twp7-W4
48	<i>Balsamorhiza sagittata</i>	27-06-90	S17-R29-Twp7-W4
53	<i>Balsamorhiza sagittata</i>	27-06-90	S16-R29-Twp7-W4

**Appendix IV.** Native seed collection (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
59	<i>Balsamorhiza sagittata</i>	28-06-90	S16-R29-Twp7-W4
85	<i>Balsamorhiza sagittata</i>	10-07-90	S17-R29-Twp7-W4
113	<i>Besseyia wyomingensis</i>	08-08-90	S19-R29-Twp7-W4
134	<i>Campanula rotundifolia</i>	10-08-90	S16-R29-Twp7-W4
114	<i>Castilleja lutescens</i>	08-08-90	S19-R29-Twp7-W4
108	<i>Cirsium undulatum</i>	09-08-90	S32-R1-Twp7-W5
155	<i>Clematis ligusticifolia</i>	27-08-90	S16-R29-Twp7-W4
125	<i>Danthonia parryi</i>	09-08-90	S32-R1-Twp7-W5
46	<i>Dodecatheon conjugens</i>	27-06-90	S34-R30-Twp7-W4
82	<i>Eriogonum flavum</i>	19-07-90	S16-R29-Twp7-W4
163	<i>Festuca campestris</i>	28-06-90	S16-R29-Twp7-W4
52	<i>Fritillaria pudica</i>	27-06-90	S16-R29-Twp7-W4
54	<i>Fritillaria pudica</i>	27-06-90	S17-R29-Twp7-W4
69	<i>Fritillaria pudica</i>	10-07-90	S16-R29-Twp7-W4
105	<i>Gaillardia aristata</i>	09-08-90	S32-R1-Twp7-W5
152	<i>Gaillardia aristata</i>	27-08-90	S17-R29-Twp7-W4
137	<i>Galium boreale</i>	27-08-90	S16-R29-Twp7-W4
49	<i>Geum triflorum</i>	27-06-90	S17-R29-Twp7-W4
58	<i>Geum triflorum</i>	28-06-90	S33-R1-Twp7-W5
148	<i>Glycyrrhiza lepidota</i>	27-08-90	S16-R29-Twp7-W4
157	<i>Haplopappus spinulosus</i>	27-08-90	S17-R29-Twp7-W4
141	<i>Hedysarum boreale</i>	27-08-90	S16-R29-Twp7-W4
156	<i>Helianthus annuus</i>	27-08-90	S17-R29-Twp7-W4
143	<i>Helianthus nuttallii</i>	27-08-90	S16-R29-Twp7-W4
144	<i>Heterotheca villosa</i>	27-08-90	S17-R29-Twp7-W4
107	<i>Heuchera parvifolia</i>	08-08-90	S19-R29-Twp7-W4
99	<i>Heuchera parvifolia</i>	09-08-90	S32-R1-Twp7-W5



**Appendix IV.** Native seed collection (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
71	<i>Hymenoxys acaulis</i>	21-07-90	S17-R29-Twp7-W4
166	<i>Koeleria macrantha</i>	28-06-90	S33-R1-Twp7-W5
138	<i>Liatris punctata</i>	27-08-90	S17-R29-Twp7-W4
170	<i>Liatris punctata</i>	26-09-90	S19-R29-Twp7-W4
172	<i>Liatris punctata</i>	26-09-90	S16-R29-Twp7-W4
77	<i>Linum lewisii</i>	19-07-90	S16-R29-Twp7-W4
92	<i>Linum lewisii</i>	19-07-90	S16-R29-Twp7-W4
75	<i>Linum lewisii</i>	21-07-90	S17-R29-Twp7-W4
116	<i>Linum lewisii</i>	08-08-90	S19-R29-Twp7-W4
121	<i>Linum lewisii</i>	08-08-90	S11-R1-Twp7-W5
70	<i>Linum rigidum</i>	21-07-90	S17-R29-Twp7-W4
123	<i>Linum rigidum</i>	09-08-90	S32-R1-Twp7-W5
50	<i>Lithospermum ruderales</i>	27-06-90	S16-R29-Twp7-W4
57	<i>Lithospermum ruderales</i>	28-06-90	S16-R29-Twp7-W4
94	<i>Lithospermum ruderales</i>	10-07-90	S17-R29-Twp7-W4
115	<i>Lithospermum ruderales</i>	08-08-90	S19-R29-Twp7-W4
51	<i>Lomatium triternatum</i>	27-06-90	S34-R30-Twp7-W4
95	<i>Lomatium triternatum</i>	10-07-90	S17-R29-Twp7-W4
56	<i>Lupinus argenteus</i>	27-06-90	S16-R29-Twp7-W4
91	<i>Lupinus argenteus</i>	10-07-90	S16-R29-Twp7-W4
73	<i>Lupinus argenteus</i>	19-07-90	S16-R29-Twp7-W4
83	<i>Lupinus argenteus</i> (white)	19-07-90	S16-R29-Twp7-W4
74	<i>Lupinus sericeus?</i>	10-07-90	S16-R29-Twp7-W4
80	<i>Lupinus sericeus?</i>	19-07-90	S16-R29-Twp7-W4
68	<i>Lupinus sericeus?</i>	20-07-90	S19-R29-Twp7-W4
66	<i>Lupinus sericeus?</i>	21-07-90	S17-R29-Twp7-W4

**Appendix IV.** Native seed collection (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
124	<i>Lupinus sericeus?</i>	08-08-90	S19-R29-Twp7-W4
104	<i>Lupinus sericeus?</i>	09-08-90	S32-R1-Twp7-W5
110	<i>Lupinus sericeus?</i>	09-08-90	S32-R1-Twp7-W5
142	<i>Monarda fistulosa</i>	27-08-90	S16-R29-Twp7-W4
154	<i>Monarda fistulosa</i>	27-08-90	S16-R29-Twp7-W4
112	<i>Orobanche fasciculata</i>	08-08-90	S19-R29-Twp7-W4
81	<i>Oryzopsis hymenoides</i>	19-07-90	S16-R29-Twp7-W4
119	<i>Oxytropis monticola</i>	08-08-90	S19-R29-Twp7-W4
182	<i>Oxytropis sericea</i>	03-09-90	S33-R1-Twp7-W5
72	<i>Oxytropis sericea</i>	10-07-90	S16-R29-Twp7-W4
78	<i>Oxytropis sericea</i>	19-07-90	S16-R29-Twp7-W4
102	<i>Oxytropis sericea</i>	08-08-90	S19-R29-Twp7-W4
135	<i>Oxytropis sericea</i>	10-08-90	S16-R29-Twp7-W4
103	<i>Penstemon confertus</i>	09-08-90	S32-R1-Twp7-W5
173	<i>Penstemon nitidus</i>	09-08-90	S16-R29-Twp7-W4
130	<i>Penstemon nitidus</i>	10-08-90	S16-R29-Twp7-W4
145	<i>Petalostemon candidum</i>	27-08-90	S16-R29-Twp7-W4
93	<i>Potentilla hippiana</i>	19-07-90	S16-R29-Twp7-W4
63	<i>Potentilla hippiana</i>	21-07-90	S17-R29-Twp7-W4
106	<i>Potentilla hippiana</i>	08-08-90	S11-R1-Twp7-W5
111	<i>Potentilla pensylvanica</i>	08-08-90	S19-R29-Twp7-W4
151	<i>Potentilla</i> sp.	27-08-90	S16-R29-Twp7-W4
127	<i>Psoralea esculenta</i>	09-08-90	S17-R29-Twp7-W4
136	<i>Psoralea esculenta</i>	10-08-90	S16-R29-Twp7-W4
147	<i>Psoralea esculenta</i>	27-08-90	S16-R29-Twp7-W4
146	<i>Ratibida columnifera</i>	27-08-90	S16-R29-Twp7-W4
171	<i>Ratibida columnifera</i>	26-09-90	S16-R29-Twp7-W4

**Appendix IV.** Native seed collection (some identifications have yet to be verified).

Coll. No.	Taxa	Date	Location
84	<i>Senecio canus</i>	10-07-90	S16-R29-Twp7-W4
117	<i>Silene noctiflora</i>	08-08-90	S19-R29-Twp7-W4
96	<i>Sisyrinchium montanum</i>	19-07-90	S16-R29-Twp7-W4
133	<i>Sisyrinchium montanum</i>	10-08-90	S16-R29-Twp7-W4
109	<i>Stipa curtisetia</i>	09-08-90	S32-R1-Twp7-W5
129	<i>Stipa</i> sp.	10-08-90	S16-R29-Twp7-W4
126	<i>Stipa viridula</i>	31-08-90	S33-R1-Twp7-W5
64	<i>Thermopsis rhombifolia</i>	21-07-90	S24-R1-Twp7-W5
160	<i>Viola nuttallii</i>	21-06-90	S17-R29-Twp7-W4
101	<i>Zigadenus venenosus</i>	09-08-90	S32-R1-Twp7-W5

Appendix IV. Native seed collection from identification tags set in  
at 100m intervals along the riverbank. (See also Appendix I)

Col.	Loc.	Native seed	Location	Notes
101	101	101	101	101
102	102	102	102	102
103	103	103	103	103
104	104	104	104	104
105	105	105	105	105
106	106	106	106	106
107	107	107	107	107
108	108	108	108	108
109	109	109	109	109
110	110	110	110	110
111	111	111	111	111
112	112	112	112	112
113	113	113	113	113
114	114	114	114	114
115	115	115	115	115
116	116	116	116	116
117	117	117	117	117
118	118	118	118	118
119	119	119	119	119
120	120	120	120	120
121	121	121	121	121
122	122	122	122	122
123	123	123	123	123
124	124	124	124	124
125	125	125	125	125
126	126	126	126	126
127	127	127	127	127
128	128	128	128	128
129	129	129	129	129
130	130	130	130	130
131	131	131	131	131
132	132	132	132	132
133	133	133	133	133
134	134	134	134	134
135	135	135	135	135
136	136	136	136	136
137	137	137	137	137
138	138	138	138	138
139	139	139	139	139
140	140	140	140	140
141	141	141	141	141
142	142	142	142	142
143	143	143	143	143
144	144	144	144	144
145	145	145	145	145
146	146	146	146	146
147	147	147	147	147
148	148	148	148	148
149	149	149	149	149
150	150	150	150	150





N.L.C. - B.N.C.



3 3286 11067119 1